

ISO 19290:2021 (E)

Cigarettes — Determination of tobacco specific nitrosamines in mainstream cigarette smoke — Method using LC-MS/MS

Contents

	Foreword
	Introduction
1	Scope
2	Normative references
3	Terms and definitions
4	Principle
5	Apparatus
6	Reagents
7	Preparation
7.1	Preparation of glassware
7.2	Preparation of solutions
7.3	Preparation of standards
7.3.1	General
7.3.2	Preparation of internal standard solutions
7.3.2.1	Primary solution
7.3.2.2	Combined secondary solution
7.3.2.3	Working solution
7.3.3	Preparation of calibration standard solutions
7.3.3.1	Primary single TSNA solutions
7.3.3.2	Mixed TSNA stock solution (I)
7.3.3.3	Mixed TSNA stock solution (II)
7.3.3.4	Working standard solutions
7.3.3.5	Storage
8	Sampling
9	Tobacco product preparation
10	Sample generation — Smoking of cigarettes
10.1	General
10.2	Linear smoking
10.3	Rotary smoking
11	Sample analysis
11.1	Preparation of sample
11.1.1	General
11.1.2	Extraction for linear smoking (44 mm pad)
11.1.3	Extraction for rotary smoking (92 mm pad)
11.1.4	Final sample preparation
11.2	Reversed phase high performance liquid chromatography
11.2.1	General
11.2.2	HPLC set-up parameters (Example)
11.2.3	Mobile phase (Example)
11.2.4	Mobile phase: Gradient (Example)
11.2.5	MS/MS Set-up parameters (Example)

11.3	Calculation
11.3.1	Calibration curve
11.3.2	Determination of the TSNA concentrations
11.3.3	Sample quantification
11.3.4	Determination of mainstream smoke TSNA deliveries
12	Repeatability and reproducibility
13	Test report
Annex A	(informative) Example of typical chromatograms

Page count: 14